

Equivariant Twisted Cartan Cohomology Theory

Debasis Sen

Abstract

In this note we prove an equivariant version of a result of Cartan [Car76] for equivariant simplicial cohomology with local coefficients.

Keywords: Simplicial sets, group action, local coefficients, Cartan cohomology theory, equivariant twisted cohomology, generalized Eilenberg-MacLane complex.

1 Introduction

To generalize Sullivan's theory of rational de Rham complexes on simplicial sets to cochain complexes over arbitrary ring of coefficients, Cartan [Car76] introduced the notion of 'Cohomology theory'. Over the coefficient ring $\mathbb{Z}$, Cartan's result can be described as follows. Recall that a *simplicial differential graded algebra* over $\mathbb{Z}$ is a simplicial object in the category DGA of differential graded algebras over $\mathbb{Z}$, so that for each $p \geq 0$ we have a differential graded algebra

$$(A_p^*, \delta): A_p^0 \xrightarrow{\delta} A_p^1 \xrightarrow{\delta} A_p^2 \rightarrow \dots$$

together with face and degeneracy maps $\partial_i: A_{p+1}^* \rightarrow A_p^*$ and $s_i: A_p^* \rightarrow A_{p+1}^*$ which are homomorphism of differential graded algebras satisfying the usual simplicial and differential identities. Then a *cohomology theory* in the sense of Cartan is a simplicial differential graded algebra A over $\mathbb{Z}$ such that

1. each cochain complex (A_p^*, δ) is exact and $Z^0 A = \text{Ker}(A_*^0 \xrightarrow{\delta} A_*^1)$ is a simplicially trivial algebra over $\mathbb{Z}$ (here simplicially trivial means that all the face and degeneracy maps are isomorphisms),
2. the homotopy groups $\pi_i(A_*^n)$ of the simplicial set $A_*^n = \{A_p^n\}_{p \geq 0}$ are trivial for all $i, n \geq 0$.

A cohomology theory A determines a contravariant functor from the category of simplicial sets to DGA which assigns to each simplicial set K the differential

The author would like to thank CSIR for its support.
Mathematics Subject Classifications (2010): 55U10, 55U35, 55N91, 55N25, 57S99.

graded algebra $A(K) = \{Hom(K, A_*^n)\}_{n \geq 0}$, where $Hom(K, A_*^n)$ is the abelian group of simplicial maps $K \rightarrow A_*^n$ and the differential on $A(K)$ is induced from that of A . Then Cartan's theorem states that there is a natural isomorphism

$$H^*(A(K)) \cong H^*(K; \mathbb{Z}(A)),$$

where $\mathbb{Z}(A)$ is the abelian group $(Z^0 A)_0$.

In [Hir79], Hirashima generalized Cartan's result for cohomology with local coefficients. Moreover Cartan's theorem was generalized in [MN98] for G -simplicial sets, i.e for simplicial sets equipped with an action of a group G by simplicial maps. In the equivariant setting, ordinary cohomology of simplicial sets is replaced by Bredon cohomology of G -simplicial sets.

Recently, in [MS10] the notion of equivariant cohomology with local coefficients, called Bredon-Illman cohomology with local coefficients, has been formulated for G -simplicial sets. This is the simplicial version of the equivariant cohomology with local coefficients for G -spaces as introduced in [MM96], which generalizes Bredon-Illman cohomology [Bre67], [Ill75]. It is therefore reasonable to prove a version of Cartan's theorem on G -simplicial sets for equivariant cohomology with local coefficients. In this note we define the notion of equivariant twisted Cartan cohomology theory and prove a version of Cartan's theorem which reduces to the result of [MN98] when the local coefficients system is simple, and, to that of [Hir79] when G is a trivial group. In the present context Cartan's cohomology theory appears as a contravariant functor from the category O_G of canonical orbits to the category of cohomology theory in the sense of Cartan [Car76], satisfying some naturality conditions (see Definition 4.1 for details). It may be remarked that the equivariant analogue of a point is the orbit of the point and these orbits are precisely the objects of the category O_G .

Throughout the paper, when referring to 'model', we will always mean it to be minimal.

This paper is organized as follows. Section 2 is a review of the basic definitions and results that will be used in the sequel. In section 3, we describe the notion of equivariant twisted cohomology and state a classification theorem which is proved in [MS10]. In Section 4, we define the notion of equivariant Cartan cohomology theory and prove our main result.

2 Preliminaries

In this section we recall some basic definitions and facts about simplicial sets [May67] and related topics. We denote the category of simplicial sets and simplicial maps by $\mathcal{S}$ and the category of simplicial groups and simplicial group homomorphisms by $S\mathcal{G}$. Throughout G will denote a discrete group.

Definition 2.1. [Moore56]

Let B be a simplicial set and Γ a simplicial group. Then a graded function

$$\tau: B \longrightarrow \Gamma, \quad \tau_q: B_q \rightarrow \Gamma_{q-1}$$

is called a twisting function if it satisfies the following identities:

$$\begin{aligned}\partial_0(\tau_q(b)) &= (\tau_{q-1}(\partial_0 b))^{-1} \tau_{q-1}(\partial_1 b), \quad b \in B_q \\ \partial_i(\tau_q(b)) &= \tau_{q-1}(\partial_{i+1} b) \quad i > 0 \\ s_i(\tau_q(b)) &= \tau_{q+1}(s_{i+1} b) \quad i \geq 0 \\ \tau_{q+1}(s_0 b) &= e_q, \quad e_q \text{ being the identity of the group } \Gamma_q.\end{aligned}$$

Definition 2.2. Let B, F be simplicial sets, Γ a simplicial group which operates on F from the left, and $\tau: B \rightarrow \Gamma$ a twisting function. A twisted cartesian product (TCP), with fibre F , base B and group Γ is a simplicial set, denoted by $F \times_\tau B$ which satisfies

$$(F \times_\tau B)_n = F_n \times B_n$$

and has face and degeneracy operators

$$\begin{aligned}\partial_i(f, b) &= (\partial_i f, \partial_i b), \quad i > 0 \\ \partial_0(f, b) &= (\tau(b) \partial_0 f, \partial_0 b) \\ s_i(f, b) &= (s_i f, s_i b) \quad i \geq 0\end{aligned}$$

If B, F are Kan complexes then $F \times_\tau B$ is also a Kan complex and the canonical projection $p: F \times_\tau B \rightarrow B$ is a Kan fibration.

For an abelian group A and an integer $n > 1$, let $K(A, n)$ denote a minimal Eilenberg MacLane complex of type (A, n) . There is a canonical model of $K(A, n)$ for which the q -simplices are described as follows. Consider the simplicial abelian group $C(A, n)$ with q -simplices

$$C(A, n)_q = C^n(\Delta[q]; A),$$

the group of normalized n -cochains of the standard simplicial q -simplex $\Delta[q]$ [May67]. The face and degeneracy maps of $C(A, n)$ are given as follows. For $\mu \in C(A, n)_q$, $\alpha \in \Delta[q-1]_n$ and $\beta \in \Delta[q+1]_n$

$$\partial_i \mu(\alpha) = \mu(\delta_i(\alpha)), \quad s_j \mu(\beta) = \mu(\sigma_j(\beta)).$$

Here $\delta_i: \Delta[q-1] \rightarrow \Delta[q]$ and $\sigma_j: \Delta[q+1] \rightarrow \Delta[q]$ are the simplicial maps defined by $\delta_i(\Delta_{q-1}) = \partial_i \Delta_q$, $\sigma_j(\Delta_{q+1}) = \Delta_q$, $\Delta_q = (0, 1, \dots, q)$ being the unique non-degenerate q -simplex of $\Delta[q]$.

We have a simplicial group homomorphism

$$\delta^n: C(A, n) \rightarrow C(A, n+1)$$

such that $\delta^n c \in C(A, n+1)_q$ is the usual simplicial coboundary of $c \in C(A, n)_q$. Then

$$K(A, n)_q = \text{Ker } \delta^n = Z^n(\Delta[q]; A)$$

the group of normalized n -cocycles. It may be noted that $K(A, n)$ is a minimal one vertex Kan complex.

Definition 2.3. Let π be a group. A π -module is a pair (A, ϕ) where A is an abelian group and $\phi: \pi \rightarrow \text{Aut}(A)$ a group homomorphism. A map of π -modules $f: (A, \phi) \rightarrow (A', \phi')$ is a group homomorphism $f: A \rightarrow A'$ such that

$$f(\phi(x)a) = \phi'(x)f(a)$$

for all $x \in \pi$ and $a \in A$. The category of π -modules is denoted by $\pi\text{-mod}$.

Let $(A, \phi) \in \pi\text{-mod}$. Then π acts on the minimal one vertex Kan complex $K(A, n)$ in the following way:

$$x\mu = \phi(x) \circ \mu \text{ where } \mu \in K(A, n)_q = Z^n(\Delta[q]; A), x \in \pi.$$

The notion of a *generalized Eilenberg MacLane complex* appears in [Git63], [Hir79], [BFGM03]. Roughly speaking, a generalized Eilenberg MacLane complex is a one vertex minimal Kan complex having exactly two non-vanishing homotopy groups, one of them being the fundamental group. It appears as the total space of a Kan fibration. Gitler [Git63] used it in the construction of cohomology operations in cohomology with local coefficients. It also plays a crucial role in classifying cohomology with local coefficients [Hir79], [BFGM03]. It may be remarked that a product of Eilenberg MacLane complexes is also sometimes referred to as a generalized Eilenberg MacLane complex.

A generalized Eilenberg MacLane complex can be constructed as follows. Let $\overline{W}\pi$ denotes the standard $\overline{W}$ construction [May67] of a group π . Let (A, ϕ) be a π -module. We have a twisting function

$$\tau(\pi): \overline{W}\pi \rightarrow \pi, \text{ where } \tau(\pi)(x_1, \dots, x_q) = x_1, x_i \in \pi,$$

and π is considered as a simplicial group with each component π and all the face and the degeneracy maps are identities. For $n > 1$ let

$$L_\pi(A, n) = K(A, n) \times_{\tau(\pi)} \overline{W}\pi,$$

where the right hand side is the twisted cartesian product as defined in the Definition 2.2. Then it is a one vertex minimal Kan complex whose fundamental group is π , n -th homotopy group is A and all other homotopy groups are trivial. Moreover the action of the fundamental group π on the n -th homotopy group A is given by ϕ [Thu97]. We have a canonical map $p: L_\pi(A, n) \rightarrow \overline{W}\pi$, $p(c, x) = x$ for $c \in K, x \in \overline{W}\pi$, which is a Kan fibration.

For a group G , the *category of canonical orbits*, denoted by O_G , is a category whose objects are cosets G/H , as H runs over subgroups of G . A morphism from G/H to G/K is a G -map. Recall that such a morphism determines and is determined by a subconjugacy relation $g^{-1}Hg \subseteq K$ and is given by $\hat{g}(eH) = gK$. We denote this morphism by $\hat{g}$ [Bre67].

A contravariant functor from O_G to $\mathcal{S}$ (resp. the category of groups or the category of abelian groups) is called an O_G -*simplicial set* (resp. O_G -*group* or *abelian O_G -group*). We denote by $O_G\mathcal{S}$, the category of O_G -simplicial sets with morphisms being natural transformations of functors.

A morphism $f: T \rightarrow S$ of O_G -simplicial sets is called an O_G -Kan fibration if $f(G/H): T(G/H) \rightarrow S(G/H)$ is a Kan fibration for each subgroup H of G . Similarly, an O_G -simplicial set T is called an O_G -Kan complex if each $T(G/H)$ is a Kan complex for each subgroup $H \subseteq G$.

We recall the following definition from [MN98].

Definition 2.4. *Given an O_G -group λ and an integer $n \geq 0$, an O_G -Kan complex T is called an O_G -Eilenberg MacLane complex of type (λ, n) if each $T(G/H)$ is a $K(\lambda(G/H), n)$ and $T(\hat{g}): T(G/H) \rightarrow T(G/K)$ is the unique simplicial homomorphism induced by the linear map $\lambda(\hat{g}): \lambda(G/H) \rightarrow \lambda(G/K)$, $g^{-1}Hg \subseteq K$, such that $T(\hat{g})_n: K(\lambda(G/H), n)_n \rightarrow K(\lambda(G/K), n)_n$ is $\lambda(\hat{g})$.*

It is proved in [MN98] that any two O_G -Eilenberg MacLane complexes of the same type are naturally isomorphic. We denote an O_G -Eilenberg MacLane complex of type (λ, n) by $K(\lambda, n)$. Let $n > 1$ and λ be an abelian O_G -group. Using the canonical model of an ordinary Eilenberg MacLane complex as described at the beginning of this section, we have a canonical model of $K(\lambda, n)$, given by $K(\lambda, n)(G/H)_q = Z^n(\Delta[q]; \lambda(G/H))$.

Let $\mathcal{C}$ be a category and $T: O_G \rightarrow \mathcal{C}$ be a contravariant functor. An O_G -group $\underline{\pi}$ is said to act on T if we have a group homomorphism $\phi_H: \underline{\pi}(G/H) \rightarrow \text{Aut}_{\mathcal{C}}(T(G/H))$ for each subgroup H of G such that for any subconjugacy relation $g^{-1}Hg \subseteq K$,

$$\phi_H(\underline{\pi}(\hat{g})v) \circ T(\hat{g}) = T(\hat{g}) \circ \phi_K(v), \quad v \in \underline{\pi}(G/K).$$

We denote this action simply by ϕ . Thus we can talk of an action of $\underline{\pi}$ on an O_G -simplicial set, an O_G -group etc. If $\underline{\pi}$ acts on an abelian O_G -group T , then we call T a $\underline{\pi}$ -module.

3 G-simplicial set, Equivariant Twisted Cohomology and its Classification

In this section we briefly recall the definition of equivariant twisted cohomology and its homotopy classification from [MS10]. Let G be a discrete group. Recall that a G -simplicial set is a simplicial set $X = \{X_n\}$ such that each X_n is a G -set and the face, degeneracy maps commute with this action. A G -simplicial set X is called G -connected if each fixed point simplicial set X^H , $H \subseteq G$, is connected.

Let X be a G -simplicial set and $\underline{\pi}$ be an O_G -group. Let ΦX denote the O_G -simplicial set defined by $\Phi X(G/H) = X^H$, $\Phi X(\hat{g})(x) = gx$, $x \in X^H$, $g^{-1}Hg \subseteq K$.

Definition 3.1. *Let T be an O_G -simplicial set and Γ a simplicial O_G -group. A natural transformation of functors $\tau: T \rightarrow \Gamma$ is called an O_G -twisting function if $\tau(G/H): T(G/H) \rightarrow \Gamma(G/H)$ is an ordinary twisting function for each subgroup H of G .*

Example 3.2. Consider the O_G -group $\underline{\pi}$ as a simplicial O_G -group $\{\underline{\pi}_n\}_{n \geq 0}$ where $\underline{\pi}_n = \underline{\pi}$ for all $n \geq 0$ and face and degeneracy maps are identity natural transformations. Define

$$\tau(\underline{\pi}): \overline{W}\underline{\pi} \rightarrow \underline{\pi}, \quad \tau(\underline{\pi})(G/H)([x_1, \dots, x_q]) = x_1,$$

where $[x_1, \dots, x_q] \in \overline{W}\underline{\pi}(G/H)_q$, $x_i \in \underline{\pi}(G/H)$, $1 \leq i \leq q$. It is routine to check that $\tau(\underline{\pi})$ is an O_G -twisting function.

Example 3.3. Let X be a G -connected G -simplicial set and v be a G -fixed 0-simplex in X . Let $\underline{\pi}X: O_G \rightarrow \text{Grp}$ be the O_G -group defined as follows. For any subgroup H of G ,

$$\underline{\pi}X(G/H) = \pi_1(X^H, v)$$

and for a morphism $\hat{g}: G/H \rightarrow G/K$, $g^{-1}Hg \subseteq K$, $\underline{\pi}X(\hat{g})$ is the homomorphism of fundamental groups induced by the simplicial map $g: X^K \rightarrow X^H$. We regard $\underline{\pi}X$ as an O_G -group complex in the trivial way, that is, $\underline{\pi}X(G/H)_n = \underline{\pi}X(G/H)$ for all n . We choose a 0-simplex x on each G -orbit of X_0 and a 1-simplex $\omega_x \in X^{G_x}$ such that $\partial_0 \omega_x = x$, $\partial_1 \omega_x = v$. For any other 0-simplex y on the orbit of x we define $\omega_y = g\omega_x$ if $y = gx$. Then it is an easy check that this is well defined and $\omega_y \in X_1^{G_y}$. For a 0-simplex $x \in X^H$, let $\xi_H(x) = [\overline{\omega}_x]$ be the homotopy class of $\overline{\omega}_x: \Delta[1] \rightarrow X^H$. Here for any q -simplex σ of a simplicial set Y , $\overline{\sigma}: \Delta[q] \rightarrow Y$ denote the unique simplicial map satisfying $\overline{\sigma}(\Delta_q) = \sigma$. Define

$$\{\kappa(G/H)_n\}: X^H \rightarrow \pi_1(X^H, v)$$

by

$$\kappa(G/H)_n(y) = \xi_H(\partial_{(0,2,\dots,n)}y)^{-1} \circ [\overline{\partial_{(2,\dots,n)}y}] \circ \xi_H(\partial_{(1,\dots,n)}y)$$

where $y \in (X^H)_n$ and

$$\partial_{(0,2,\dots,n)}y = \partial_0 \partial_2 \cdots \partial_n y, \partial_{(2,\dots,n)}y = \partial_2 \cdots \partial_n y, \partial_{(1,2,\dots,n)}y = \partial_1 \partial_2 \cdots \partial_n y.$$

It is standard that $\kappa(G/H)$ is a twisting function on X^H . We verify that

$$\kappa: \Phi X \rightarrow \underline{\pi}X, \quad G/H \mapsto \kappa(G/H)$$

is natural. Suppose H and K are subgroups such that $g^{-1}Hg \subseteq K$. Let $z \in X_n^K$. Then $y = gz \in X_n^H$. Observe that if $x_1, x_2 \in X_1^K$ are 1-simplexes such that $\overline{x}_1 \simeq \overline{x}_2$, as simplicial maps into X^K then $\overline{y}_1 \simeq \overline{y}_2$ as simplicial maps into X^H where $y_i = gx_i$, $i = 1, 2$. Thus

$$\begin{aligned} \kappa(G/H)_n \circ \Phi X(\hat{g})(z) &= \kappa(G/H)_n(y) \\ &= \xi_H(\partial_{(0,2,\dots,n)}y)^{-1} \circ [\overline{\partial_{(2,\dots,n)}y}] \circ \xi_H(\partial_{(1,\dots,n)}y) \\ &= \xi_H(g\partial_{(0,2,\dots,n)}z)^{-1} \circ [\overline{g\partial_{(2,\dots,n)}z}] \circ \xi_H(g\partial_{(1,\dots,n)}z) \\ &= g\xi_K(\partial_{(0,2,\dots,n)}z)^{-1} \circ g[\overline{\partial_{(2,\dots,n)}z}] \circ g\xi_K(\partial_{(1,\dots,n)}z) \\ &= \underline{\pi}X(\hat{g}) \circ \kappa(G/K)_n(z). \end{aligned}$$

Thus $\kappa: \Phi X \rightarrow \underline{\pi}X$ is an O_G -twisting function.

Let X be a G -simplicial set, $\tau: \Phi X \rightarrow \underline{\pi}$ be an O_G -twisting function and M a $\underline{\pi}$ -module, given by ϕ . We define equivariant twisted cohomology of (X, τ, ϕ) as follows.

We denote the category of abelian O_G -groups by $\mathcal{C}_G$. We have a cochain complex in the abelian category $\mathcal{C}_G$ defined by

$$\underline{C}_n(X): O_G \rightarrow Ab, \quad G/H \mapsto C_n(X^H; \mathbb{Z}),$$

where $C_n(X^H; \mathbb{Z})$ is the free abelian group generated by the non-degenerate n -simplexes of X^H and for any morphism $\hat{g}: G/H \rightarrow G/K$, $g^{-1}Hg \subseteq K$ in O_G , $\underline{C}_n(X)(\hat{g})$ is given by the map $g_*: C_n(X^K; \mathbb{Z}) \rightarrow C_n(X^H; \mathbb{Z})$, induced by the simplicial map $g: X^K \rightarrow X^H$. The boundary $\partial_n: \underline{C}_n(X) \rightarrow \underline{C}_{n-1}(X)$ is a natural transformation defined by $\partial_n(G/H): C_n(X^H; \mathbb{Z}) \rightarrow C_{n-1}(X^H; \mathbb{Z})$, where $\partial_n(G/H)$ is the ordinary boundary map of the simplicial set X^H . Dualising this chain complex in the abelian category $\mathcal{C}_G$ we get the cochain complex

$$\{C_G^*(X; M) = \text{Hom}_{\mathcal{C}_G}(\underline{C}_*(X), M), \delta^n\},$$

which defines the ordinary Bredon cohomology of the G -simplicial set X with coefficients M [Bre67]. To define the twisted cohomology of the G -simplicial set X we modify the coboundary maps as follows

$$\delta_\tau^n: C_G^n(X; M) \rightarrow C_G^{n+1}(X; M), \quad f \mapsto \delta_\tau^n f$$

where

$$\delta_\tau^n f(G/H): C_{n+1}(X^H; \mathbb{Z}) \rightarrow M(G/H)$$

is given by

$$\delta_\tau^n f(G/H)(x) = (\tau(G/H)_{n+1}(x))^{-1} f(G/H)(\partial_0 x) + \sum_{i=1}^{n+1} (-1)^i f(G/H)(\partial_i x)$$

for $x \in X_{n+1}^H$. Note that the first term of the right hand side is obtained by the given action ϕ . We denote the resulting cochain complex by $C_G^*(X; \tau, \phi)$.

Definition 3.4. *The n^{th} equivariant twisted cohomology of (X, τ, ϕ) is defined as*

$$H_G^n(X; \tau, \phi) = H_n(C_G^*(X; \tau, \phi)).$$

Suppose that B, F are O_G -Kan complexes and Γ an O_G -group complex. Also assume that B is a Γ -module and $\kappa: B \rightarrow \Gamma$ an O_G -twisting function. Then we have the O_G -Kan complex $F \times_\kappa B$, defined as

$$(F \times_\tau B)(G/H) = F(G/H) \times_{\tau(G/H)} B(G/H), \quad (F \times_\tau B)(\hat{g}) = (F(\hat{g}), B(\hat{g})),$$

for each object G/H and morphism $\hat{g}: G/H \rightarrow G/K$ of the category O_G . We call this O_G -Kan complex the *O_G -twisted cartesian product* (TCP), with fibre F , base B , group Γ and twisting κ . Observe that the second factor projection gives an O_G -Kan fibration $p: (F \times_\tau B) \rightarrow B$. We view $(F \times_\tau B, p)$ as an object in the slice category (cf. [GJ99]) $O_G \mathcal{S}/B$.

Let M be a $\underline{\pi}$ -module with module structure given by ϕ . For each subgroup H of G , define a group homomorphism

$$\psi_H: \underline{\pi}(G/H) \rightarrow \text{Aut}_{\mathcal{S}}(K(M(G/H), n))$$

as follows. For $u \in \underline{\pi}(G/H)$, let $\psi_H(u)$ be the unique simplicial automorphism of $K(M(G/H), n)$ such that

$$\phi_H(u) = \psi_H(u)_n: K(M(G/H)_n) \rightarrow K(M(G/H)_n), \quad u \in \underline{\pi}(G/H).$$

This defines an action of the O_G -group $\underline{\pi}$ on the O_G -Kan complex $K(M, n)$. Therefore we can form the O_G -Kan fibration $p: K(M, n) \times_{\tau(\underline{\pi})} \overline{W}\underline{\pi} \rightarrow \overline{W}\underline{\pi}$, where $\tau(\underline{\pi})$ is the O_G -twisting function as described in the Example 3.2. If we use the canonical model of $K(M, n)$, the total complex of the resulting O_G -Kan fibration is denoted by $L_\phi(M, n)$. Since any two models of $K(M, n)$ are naturally isomorphic, $K(M, n) \times_{\tau(\underline{\pi})} \overline{W}\underline{\pi}$ is isomorphic to $L_\phi(M, n)$ for any model of $K(M, n)$. We call $L_\phi(M, n)$ a *generalized O_G -Eilenberg MacLane complex*. Note that $L_\phi(M, n)(G/H)$ is the generalized Eilenberg MacLane complex

$$L_{\underline{\pi}(G/H)}(M(G/H), n) = Z^n(\Delta[-]; M(G/H)) \times_{\tau(\underline{\pi}(G/H))} \overline{W}\underline{\pi}(G/H).$$

The equivariant twisted cohomology $H_G^*(X; \tau, \phi)$ has been classified by the O_G -Kan complex $L_\phi(M, n)$ in [MS10]. This classification result can be described as follows.

Let X be a G -simplicial set and $\tau: \Phi X \rightarrow \overline{W}\underline{\pi}$ be an O_G -twisting function. It determines an O_G -simplicial map $\theta(\tau): \Phi X \rightarrow \overline{W}\underline{\pi}$ defined as,

$$\theta(\tau)(G/H): X_q^H \rightarrow \overline{W}\underline{\pi}(G/H)_q,$$

$$x \mapsto [\tau(G/H)_q(x), \tau(G/H)_{q-1}(\partial_0 x), \dots, \tau(G/H)_1(\partial_0^{q-1} x)].$$

Let $(\Phi X, L_\phi(M, n))_{\overline{W}\underline{\pi}}$ denote the set of liftings of the map $\theta(\tau)$ with respect to $p: L_\phi(M, n) \rightarrow \overline{W}\underline{\pi}$, $p(c, g) = g$.

Definition 3.5. Let $f, g \in (\Phi X, L_\phi(M, n))_{\overline{W}\underline{\pi}}$. Then f and g are said to be vertically homotopic, written $f \sim_v g$, if there is a map $F: \Phi X \times \Delta[1] \rightarrow L_\phi(M, n)$ of O_G -simplicial sets such that for every object G/H of O_G , $F(G/H)$ is a homotopy of the simplicial maps $f(G/H), g(G/H)$ and $p \circ F = \theta(\tau) \circ pr_1$, where $pr_1: \Phi X \times \Delta[1] \rightarrow \Phi X$ is the projection onto the first factor.

Observe that $(\Phi X, \theta(\tau))$ and $(L_\phi(M, n), p)$ are objects in the slice category $O_G\mathcal{S}/\overline{W}\underline{\pi}$ and $(\Phi X, L_\phi(M, n))_{\overline{W}\underline{\pi}}$ is the set of morphisms in $O_G\mathcal{S}/\overline{W}\underline{\pi}$ from $(\Phi X, \theta(\tau))$ to $(L_\phi(M, n), p)$. The category $O_G\mathcal{S}$ is a closed model category [DK83] in the sense of Quillen [Qui67] and hence $O_G\mathcal{S}/\overline{W}\underline{\pi}$ is also a closed model category [GJ99], where $(L_\phi(M, n), p)$ is a fibrant object and the above notion of vertical homotopy coincides with the abstract homotopy. Therefore $\sim_v$ is an equivalence relation on the set $(\Phi X, L_\phi(M, n))_{\overline{W}\underline{\pi}}$. Let $[\Phi X, L_\phi(M, n)]_{\overline{W}\underline{\pi}}$ denote the set of equivalence classes. Then the homotopy classification of equivariant twisted cohomology can be stated as follows.

Theorem 3.6. *Suppose X is a G -simplicial set and $\tau: \Phi X \rightarrow \underline{\pi}$ is an O_G -twisting function. Then*

$$H_G^n(X; \tau, \phi) \cong [\Phi X, L_\phi(M, n)]_{\overline{W}\underline{\pi}}, \text{ for each } n \geq 0.$$

4 Equivariant Twisted Cartan Cohomology Theory

In this final section we formulate an equivariant version of Cartan's Cohomology theory [Car76] and prove that Bredon-Illman cohomology with local coefficients of a G -simplicial set can be computed by the cohomology of a differential graded algebra determined by a given cohomology theory.

We begin with the following equivariant generalization of Cartan Cohomology theory suitable for our purpose.

Definition 4.1. *An equivariant twisted Cartan cohomology theory is a sequence $\mathcal{A} = \{A^i\}_{i \geq 0}$ of simplicial abelian O_G -groups A^i , together with simplicial differentials $\delta^i: A^i \rightarrow A^{i+1}$ such that*

1. *For each subgroup $H \subseteq G$, $\mathcal{A}(G/H) = (A^*(G/H)_*, \delta^*(G/H))$ is a simplicial differential graded algebra over $\mathbb{Z}$.*
2. *For each $p \geq 0$,*

$$A_p^0 \xrightarrow{\delta_p^0} A_p^1 \xrightarrow{\delta_p^1} A_p^2 \rightarrow \dots$$

is an exact sequence in the abelian category $\mathcal{C}_G$ of abelian O_G -groups.

3. *The O_G -group $\pi_n \circ A^i$ is the zero O_G -group, for all $n, i \geq 0$.*
4. *The simplicial abelian O_G -group $Z^0 \mathcal{A} = \ker(A^0 \xrightarrow{\delta^0} A^1)$ is simplicially trivial.*
5. *For each subgroup $H \subseteq G$ and an integer $i \geq 0$ there is a group homomorphism*

$$\psi_H^i: \text{Aut}((Z^0 \mathcal{A})_0(G/H)) \rightarrow \text{Aut}_{SG}(A^i(G/H))$$

satisfying

- $\delta^i \circ \psi_H^i(\alpha) = \psi_H^{i+1}(\alpha) \circ \delta^i, \alpha \in \text{Aut}((Z^0 \mathcal{A})_0(G/H)) \ i \geq 0.$
- *If $g^{-1}Hg \subseteq K$, $\alpha \in \text{Aut}((Z^0 \mathcal{A})_0(G/H))$, $\beta \in \text{Aut}((Z^0 \mathcal{A})_0(G/K))$ such that $\alpha \circ (Z^0 \mathcal{A})_0(\hat{g}) = (Z^0 \mathcal{A})_0(\hat{g}) \circ \beta$ then*

$$\psi_H(\alpha) \circ A^i(\hat{g}) = A^i(\hat{g}) \circ \psi_K(\beta).$$

Example 4.2. *For an abelian group B and an integer $n \geq 0$, let $C(B, n)$ denote the simplicial abelian group and $\delta^n: C(B, n) \rightarrow C(B, n+1)$ be the simplicial homomorphism as introduced in the Section 1. Then, for an abelian O_G -group M , $\mathcal{A} = \{A^i\}_{i \geq 0}$ where $A^n(G/H) = C(M(G/H), n)$ together with the differential δ^n , defines an equivariant twisted Cartan cohomology theory such that $(Z^0 \mathcal{A})_0 = M$.*

Lemma 4.3. *Let $\mathcal{A}: A^0 \xrightarrow{\delta} A^1 \xrightarrow{\delta} \dots$ be an equivariant twisted Cartan cohomology theory. Then each A^n is contractible as an object of $O_G\mathcal{S}$.*

Proof. Consider the abelian O_G -simplicial group $Z^n\mathcal{A}$ defined by $Z^n\mathcal{A}(G/H) = \text{Ker}(\delta^n(G/H): A^n(G/H) \rightarrow A^{n+1}(G/H))$, $Z^n\mathcal{A}(\hat{g}) = A^n(\hat{g})|_{Z^n\mathcal{A}(G/H)}$. For an integer $n \geq 0$ and a subgroup H of G , we have a short exact sequence

$$0 \rightarrow Z^n\mathcal{A}(G/H) \rightarrow A^n(G/H) \rightarrow Z^{n+1}\mathcal{A}(G/H) \rightarrow 0$$

of simplicial abelian groups. Therefore $A^n(G/H) \rightarrow Z^{n+1}\mathcal{A}(G/H)$ is a principal fibration with fibre $Z^n\mathcal{A}(G/H)$ in the category of simplicial sets, and hence a principal twisted cartesian product (PTCP) of type (W) with group complex $Z^n\mathcal{A}(G/H)$ [May67]. This PTCP of type (W) is naturally isomorphic to the universal PTCP of type (W), $W(Z^n\mathcal{A}(G/H)) \rightarrow \overline{W}(Z^n\mathcal{A}(G/H))$. But $W(Z^n\mathcal{A}(G/H))$ is contractible. The functions

$$h_{q-i}^H: W(Z^n\mathcal{A}(G/H))_q \rightarrow W(Z^n\mathcal{A}(G/H))_{q+1}, \quad 0 \leq i \leq q, \quad q \geq 0,$$

$$h_{q-i}^H(x_q, \dots, x_0) = (0_{q+1}^H, \dots, 0_{i+1}^H, \partial_0^{q-i} x_q \cdots \partial_0 x_{i+1} \cdot x_i, x_{i-1}, \dots, x_0),$$

where $x_j \in Z^n\mathcal{A}(G/H)_j$, $0 \leq j \leq q$ and 0_{q+1-r}^H is the zero elements of the abelian group $Z^n\mathcal{A}(G/H)_{q+1-r}$, $0 \leq r \leq q-i$, defines a contraction of $W(Z^n\mathcal{A}(G/H))$ which is natural with respect to morphisms of O_G . Hence $A^n(G/H)$ is also contractible and the contraction is natural. Consequently A^n is contractible as object of $O_G\mathcal{S}$. $\square$

Consider an equivariant twisted Cartan cohomology theory $\mathcal{A} = \{A^i\}_{i \geq 0}$. It determines an abelian O_G -group $(Z^0\mathcal{A})_0$. We denote it by M . Given a G -simplicial set X , an O_G -group $\underline{\pi}$, an O_G -twisting function $\tau: \Phi X \rightarrow \underline{\pi}$, and a $\underline{\pi}$ -module structure ϕ on M , we shall construct a differential graded algebra over $\mathbb{Z}$ whose cohomology will compute the equivariant twisted cohomology of (X, ϕ, τ) .

Note that, by the second condition of the fifth axiom in the Definition 4.1, A^n becomes a $\underline{\pi}$ -module by $(\psi\phi)_H = \psi_H\phi_H: \underline{\pi}(G/H) \rightarrow \text{Aut}_{\mathcal{SG}}(A^n(G/H))$. To see this, observe that for $g^{-1}Hg \subseteq K$, $v \in \underline{\pi}(G/K)$ we have

$$\phi_H(\underline{\pi}(\hat{g})v) \circ M(\hat{g}) = M(\hat{g}) \circ \phi_K(v).$$

Therefore taking $\alpha = \phi_H(\underline{\pi}(\hat{g})v)$, $\beta = \phi_K(v)$ in the second condition of the fifth axiom in the Definition 4.1, we get

$$\psi_H\phi_H(\underline{\pi}(\hat{g})v) \circ A^n(\hat{g}) = A^n(\hat{g}) \circ \psi_K\phi_K(v).$$

Consider the O_G -twisting function as introduced in the Example 3.2. We form the O_G -Kan fibration $p: A^n \times_{\tau(\underline{\pi})} \overline{W}\underline{\pi} \rightarrow \overline{W}\underline{\pi}$ by taking the O_G -twisted cartesian product as described in Section 3.

The O_G -twisting function $\tau: \Phi X \rightarrow \overline{W}\underline{\pi}$ determines a map $\theta(\tau): \Phi X \rightarrow \overline{W}\underline{\pi}$ defined by

$$\theta(\tau)(G/H)_q(x) = [\tau(G/H)(x), \tau(G/H)(\partial_0 x), \dots, \tau(G/H)(\partial_0^{q-1} x)], \quad x \in X_q^H.$$

Let $A_\phi^n(X; \tau) = \{f: \Phi X \rightarrow A^n \times_{\tau(\underline{x})} \overline{W}\underline{\pi} \mid pf = \theta(\tau)\}$. This set has an abelian group structure by fibrewise addition, fibrewise inversion and the zero section. We define a differential $\bar{\delta}^n: A_\phi^n(X; \tau) \rightarrow A_\phi^{n+1}(X; \tau)$ by

$$(\bar{\delta}^n f)(G/H)(x) = (\delta^n(G/H)c, b), \quad f \in A_\phi^n(X; \tau), x \in X^H, f(x) = (c, b).$$

It is straightforward to check that $\{A_\phi^n(X; \tau), \bar{\delta}\}$ is a cochain complex. Furthermore $A_\phi^*(X; \tau)$ admits a graded algebra structure induced from the differential graded algebra $\mathcal{A}$. The zero element of this algebra is given by the trivial lift $\mathbf{0}$, defined by

$$\mathbf{0}(G/H)_q(x) = (0_q^H, \theta(\tau)(G/H)_q(x)),$$

where $x \in X_q^H$ and 0_q^H is the zero of the abelian group $A(G/H)_q$. As before we use the notation $[\Phi X, Z^n \mathcal{A} \times_{\tau(\underline{x})} \overline{W}\underline{\pi}]_{\overline{W}\underline{\pi}}$ to denote the set of vertical homotopy classes of liftings of $\theta(\tau)$.

Proposition 4.4. *With the above notations, we have*

$$H^n(A_\phi^*(X; \tau)) = [\Phi X, Z^n \mathcal{A} \times_{\tau(\underline{x})} \overline{W}\underline{\pi}]_{\overline{W}\underline{\pi}}.$$

Proof. Clearly $\text{Ker}(\bar{\delta}^n) = (\Phi X, Z^n \mathcal{A} \times_{\tau(\underline{x})} \overline{W}\underline{\pi})_{\overline{W}\underline{\pi}}$. We now show that

$$\text{Im}(\bar{\delta}^{n-1}) = \{f \in (\Phi X, Z^n \mathcal{A} \times_{\tau(\underline{x})} \overline{W}\underline{\pi})_{\overline{W}\underline{\pi}} \mid f \sim_v \mathbf{0}\}.$$

Let $F: f \sim_v \mathbf{0}$. Consider the following left lifting problem in the closed model category $O_G\mathcal{S}/\overline{W}\underline{\pi}$ ([DK83], [GJ99]).

$$\begin{array}{ccc} \Phi X & \xrightarrow{(0, \theta(\tau)x)} & A^{n-1} \times_{\tau(\underline{x})} \overline{W}\underline{\pi} \\ \downarrow i_1 & \nearrow \tilde{F} & \downarrow \bar{\delta}^{n-1} \\ \Phi X \times \Delta[1] & \xrightarrow{F} & Z^n \mathcal{A} \times_{\tau(\underline{x})} \overline{W}\underline{\pi} \end{array}$$

Here the O_G -simplicial set $\Phi X \times \Delta[n]$ is defined by

$$(\Phi X \times \Delta[n])(G/H) = X^H \times \Delta[n], \quad (\Phi X \times \Delta[n])(\hat{g}) = (g, id), n \geq 0.$$

We identify ΦX with $\Phi X \times \Delta[0]$. The canonical inclusions $\delta_0, \delta_1: \Delta[0] \rightarrow \Delta[1]$ (see Section 1) induce natural inclusions $i_0, i_1: \Phi X \rightarrow \Phi X \times \Delta[1]$. Note that i_1 is a trivial cofibration and $\bar{\delta}^{n-1}$ is a fibration in $O_G\mathcal{S}/\overline{W}\underline{\pi}$. Hence the above left lifting problem has a solution $\tilde{F}$. Then $\tilde{F}i_0 \in A_\phi^{n-1}(X; \tau)$ such that $\bar{\delta}^{n-1}(\tilde{F}i_0) = f$. Therefore $f \in \text{Im}(\bar{\delta}^{n-1})$.

On the other hand, suppose that $f = \bar{\delta}^{n-1}h$ for $f \in A_\phi^n(X; \tau)$ and $h \in A_\phi^{n-1}(X; \tau)$. Then clearly $f \in (\Phi X, Z^n \mathcal{A} \times_{\tau(\underline{x})} \overline{W}\underline{\pi})_{\overline{W}\underline{\pi}}$. Composing h with

first factor projection map, we get a map $h': \Phi X \rightarrow A^{n-1}$ of $O_G\mathcal{S}$. But by the Lemma 4.3 A^{n-1} is contractible. Let $H: \Phi X \times \Delta[1] \rightarrow A^{n-1}$ be a contracting homotopy for the O_G -simplicial set A^{n-1} . Then define $\tilde{H}: \Phi X \times \Delta[1] \rightarrow A_\phi^{n-1}(X; \tau)$ by $\tilde{H}(x, t) = (H(x, t), \theta(\tau)x)$. Clearly $\tilde{H}: h \sim_v \mathbf{0}$ in $O_G\mathcal{S}/\overline{W}\underline{\pi}$. Hence $\overline{\delta}^{n-1} \circ \tilde{H}: f \sim_v \mathbf{0}$. This proves the proposition for $n > 0$.

For $n = 0$, we note that $H^0(A_\phi^*(X; \tau)) = (\Phi X, Z^0\mathcal{A} \times_{\tau(\underline{\pi})} \overline{W}\underline{\pi})_{\overline{W}\underline{\pi}}$ and two elements in the right hand side are homotopic if and only if they are equal. $\square$

Observe that the fourth axiom of Definition 4.1 implies $Z^0\mathcal{A}$ is an O_G -Eilenberg MacLane complex of type $(M, 0)$ and hence by induction $Z^n\mathcal{A}$ is an O_G -Eilenberg MacLane complex of type (M, n) . To justify this, consider the fibration

$$A^n(G/H) \rightarrow Z^{n+1}\mathcal{A}(G/H)$$

with fiber $Z^n\mathcal{A}(G/H)$, $H \leq G$. As noted in the Lemma 4.3, this is a PTCP with fibre $Z^n\mathcal{A}(G/H)$. Therefore if $Z^n\mathcal{A}(G/H)$ is minimal then so is $Z^{n+1}\mathcal{A}(G/H)$. But $Z^0\mathcal{A}(G/H)$, being simplicially trivial, is minimal. Hence by induction it follows that $Z^n\mathcal{A}(G/H)$ is minimal for all n . Now applying the homotopy long exact sequence to the above fibration, and using the third axiom of the Definition 4.1 together with induction on n , we see that $Z^n\mathcal{A}$ is an O_G -Eilenberg MacLane complex of type (M, n) . Hence it is isomorphic to the canonical model of $K(M, n)$. Therefore $(Z^n\mathcal{A} \times_{\tau(\underline{\pi})} \overline{W}\underline{\pi}, p)$ is isomorphic to $(L_\phi(M, n), p)$ as objects in the slice category $O_G\mathcal{S}/\overline{W}\underline{\pi}$. So we have,

$$\begin{aligned} H^n(A_\phi(X; \tau)) &= [\Phi X, Z^n\mathcal{A} \times_{\tau(\underline{\pi})} \overline{W}\underline{\pi}]_{\overline{W}\underline{\pi}} \\ &\cong [\Phi X, L_\phi(M, n)]_{\overline{W}\underline{\pi}}. \end{aligned}$$

It follows from the Theorem 3.6 that

$$H^n(A_\phi(X; \tau)) \cong H_G^n(X; \phi, \tau).$$

Thus we have proved the following theorem.

Theorem 4.5. *Suppose $\mathcal{A}$ is an equivariant twisted Cartan cohomology theory. Then for every G -simplicial set X together with an O_G -group $\underline{\pi}$, an O_G -twisting function $\tau: \Phi X \rightarrow \underline{\pi}$ and an action ϕ of $\underline{\pi}$ on the abelian O_G -group $(Z^0\mathcal{A})_0$ there is an isomorphism of graded algebras*

$$H_G^*(X; \tau, \phi) \cong H^*(\mathcal{A}(X; \tau, \phi)),$$

where $\mathcal{A}(X; \tau, \phi)$ denote the graded algebra $(A_\phi^*(X; \tau), \overline{\delta})$.

It has been shown in [MS10] that for a G -connected G -simplicial set X with a G -fixed 0-simplex, the simplicial version of Bredon Illman cohomology with local coefficients can be interpreted as an equivariant twisted cohomology for $\underline{\pi} = \underline{\pi}X$ and the O_G -twisting function κ as described in the Example 3.3 (cf. Theorem 4.7 of [MS10]). Combining it with the Theorem 4.5 we have the following result.

Theorem 4.6. *Suppose $\mathcal{A}$ is an equivariant twisted Cartan cohomology theory. Given any G -connected G -simplicial set X with a G -fixed 0-simplex and an action ϕ of πX on $(Z^0\mathcal{A})_0$, let $\mathcal{L}$ be the equivariant local coefficients (cf. Definition 3.1 and page 1020, section 3 [MS10]) determined by the πX -module $(Z^0\mathcal{A})_0$ on X . Then*

$$H_G^*(X; \mathcal{L}) \cong H^*(\mathcal{A}(X; \kappa, \phi)).$$

5 Acknowledgement

The author had many fruitful discussions with Prof. Goutam Mukherjee and would like to thank him for his useful comments.

References

- [BFGM03] M. Bullejos, E. Faro, and M. A. García-Muñoz. Homotopy colimits and cohomology with local coefficients. *Cah. Topol. Géom. Différ. Catég.*, 44(1):63–80, 2003.
- [Bre67] Glen E. Bredon. *Equivariant cohomology theories*. Lecture Notes in Mathematics, No. 34. Springer-Verlag, Berlin, 1967.
- [Car76] Henri Cartan. Théories cohomologiques. *Invent. Math.*, 35:261–271, 1976.
- [DK83] W. G. Dwyer and D. M. Kan. Function complexes for diagrams of simplicial sets. *Nederl. Akad. Wetensch. Indag. Math.*, 45(2):139–147, 1983.
- [Git63] Samuel Gitler. Cohomology operations with local coefficients. *Amer. J. Math.*, 85:156–188, 1963.
- [GJ99] Paul G. Goerss and John F. Jardine. *Simplicial homotopy theory*, volume 174 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 1999.
- [Hir79] Yasumasa Hirashima. A note on cohomology with local coefficients. *Osaka J. Math.*, 16(1):219–231, 1979.
- [Ill75] Sören Illman. Equivariant singular homology and cohomology. I. *Mem. Amer. Math. Soc.*, 1(issue 2, 156):ii+74, 1975.
- [May67] J. Peter May. *Simplicial objects in algebraic topology*. Van Nostrand Mathematical Studies, No. 11. D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto, Ont.-London, 1967.
- [MM96] Amiya Mukherjee and Goutam Mukherjee. Bredon-Illman cohomology with local coefficients. *Quart. J. Math. Oxford Ser. (2)*, 47(186):199–219, 1996.

- [MN98] Amiya Mukherjee and Aniruddha C. Naolekar. An axiomatic approach to equivariant cohomology theories. *Proc. Indian Acad. Sci. Math. Sci.*, 108(2):151–162, 1998.
- [MS10] G Mukherjee and Debasis Sen. Equivariant simplicial cohomology with local coefficients and its classification. *Topology and its Applications*, 157(1015-1032), 2010.
- [Qui67] Daniel G. Quillen. *Homotopical algebra*. Lecture Notes in Mathematics, No. 43. Springer-Verlag, Berlin, 1967.
- [Thu97] Phillip B. Thurber. Semi-localization of a one pointed Kan complex. *Pacific J. Math.*, 178(1):147–184, 1997.
- [Moore56] J. C Moore. *Seminar on algebraic homotopy theory*. Princeton, 1956(mimeographed notes).

Debasis Sen

Indian Statistical Institute, Kolkata-700108, India.

e-mail: dsen_r@isical.ac.in, sen.deba@gmail.com